


# Cosmic Rays



## Contents

### Introduction

#### Background Radiation

Earth and beyond  
Radioactivity  
 $\alpha$   $\beta$   $\gamma$  radiation

#### Cosmic Rays

100+ years of history  
Air showers

### Detectors

Geiger counter  
Cloud chamber  
Cosmic ray observatories

### Cosmic Sources

Galactic or extra-galactic  
Magnetic fields  
Ultra-high energies

### Summary

The U3A lawyers require ...



### Risk Assessment

This talk will include a number of table-top demonstrations, some of which use radioactive samples.

The samples contain naturally occurring elements and there is no risk to any members of the audience.

## Demonstrations

On the table in front of me are two types of particle detector that I will be using in live demonstrations at various points throughout this talk.



Geiger-Müller detector



Cloud Chamber

# Cosmic Rays

## Introduction

What's that clicking?

Each click signals the arrival of a sub-atomic particle passing through the detector of what is often called a Geiger counter.

Maybe it was spat out from an atom that is lurking somewhere in this room. Maybe its origins were more distant ... the result of a supernova that went off a long time ago in a galaxy far, far away.

We can't tell just from the click itself what the particle was.



## Background Radiation

*Terrestrial radioactivity and isotopes*

## Background Radiation

Radiation is all around us ... all the time.

You have lived with it all your life and it is not something to fear.

Before dealing with cosmic rays, let's get some context by considering radiation from the Earth itself.

The Earth is radioactive.

It was radioactive when it was made 4.54 billion years ago and it still is today. This is a natural consequence of the properties of some elements.




## Radioactivity

The periodic table contains many elements that are **stable** – once an atom has been made it remains essentially unchanged ... forever.

Some elements are **unstable**. They decay (change) into other elements and in the process the atomic nuclei emit sub-atomic particles.


We can visualise the particles emitted by radioactive uranium ...



# Cosmic Rays

**Demonstration**

Visualising radiation from a radioactive mineral




The rock is clearly emitting ... something. What are the particles that are leaving tracks in the detector? Where do they come from?

UNIVERSITY OF LIVERPOOL

9

**Atomic Nucleus**

An atom is a nucleus surrounded by a cloud of electrons ...




... but it is the nucleus alone that determines the radioactivity of the atom

UNIVERSITY OF LIVERPOOL

10

**Atomic Nucleus**




UNIVERSITY OF LIVERPOOL

11

**Alpha–Beta–Gamma**

The most common types of radiation are labelled  $\alpha$ ,  $\beta$  and  $\gamma$ :

- $\alpha$  Alpha rays (or alpha particles)  
 $\alpha$  = nuclei of helium atoms = 2 protons + 2 neutrons  
 Heavy particles that cannot travel far through matter
- $\beta$  Beta rays (or beta particles)  
 $\beta$  = electrons  
 Lightweight particles that can penetrate further than alphas
- $\gamma$  Gamma radiation is rather different from  $\alpha$  and  $\beta$  rays because it is a high-energy electromagnetic wave.




UNIVERSITY OF LIVERPOOL

12


# Cosmic Rays

**Demonstration**

$\alpha$  and  $\beta$  radiation from a radioactive mineral





The faint ghostly tracks are  $\alpha$  particles; the bright narrow tracks are  $\beta$  particles


 UNIVERSITY OF LIVERPOOL 13

**Radioactive Isotopes**

Isotopes are 'cousins' of the same type of atom. For instance ...

All carbon atoms have **6** protons in the nucleus (by definition).


Carbon atoms that have <b>6 neutrons</b> are the <b>stable</b> isotope C-12	Carbon atoms that have <b>8 neutrons</b> are the <b>radioactive</b> isotope C-14
	
$6 + 6 = 12$	$6 + 8 = 14$

 UNIVERSITY OF LIVERPOOL 14


**Aside: Radioactive Carbon-14**

Because C-14 atoms decay (change) into nitrogen atoms over 1000s of years they can be used in carbon dating organic material.

If they don't last long, where do the C-14 atoms come from in the first place?



They are being continuously replenished by cosmic rays hitting nuclei of nitrogen, kicking out protons and changing N-14 into C-14.

 UNIVERSITY OF LIVERPOOL 15

**Radioactive Elements**


**Potassium** (K) is the most abundant of the radioactive elements  
In your body, potassium atoms decay at the rate of 4000/sec  
It is one of largest contributions to natural radiation


**Thorium** (Th) is more abundant than uranium, and ...

**Uranium** (U) is more abundant than you think  
At 3 ppm there is 500x more uranium than gold in the Earth's crust

In the Earth's core, these radioactive elements have been emitting radiation for billions of years, heating the core and keeping it molten.


A rotating molten core is the dynamo that creates the Earth's magnetic field.



 UNIVERSITY OF LIVERPOOL 16

# Cosmic Rays

## Earth's Magnetic Field



The Earth's magnetic fields helps to protect us

## Aside: Anti-Matter

0.01% of potassium is K-40, an isotope that emits  $\beta^+$  radiation.

$\beta^+$  particles are like electrons, but with the *opposite* electrical charge. They are particles of **anti-matter**.

Hence, foods rich in potassium emit anti-matter radiation.



This gave rise to the unit of "banana equivalent dose" (BED) that compares any dose of radiation to the dose a person would be exposed to by eating one average-sized banana.

## Banana Equivalent Dose

Eating 1 banana =



Dose of cosmic rays that everybody is exposed to every day =



( double it if you fly a lot )

## Radon


Not all radioactive elements are found in solid form.

Radon is a radioactive gas formed by the radioactive decay of the small amounts of uranium that occur naturally in all rocks.

Radon concentrations vary across the UK depending on the type of ground.

For instance, radon levels are high in parts of the UK rich in granite, such as in the South West.

Radon gas atoms emit  $\alpha$  particles.



# Cosmic Rays

## Cosmic Rays

*Having considered terrestrial sources of radiation  
we can now return to the main topic of this talk*

## Pioneers

The pioneers studying radiation in the period 1895–1915 were ...



Henri Becquerel  
1852 – 1908



Marie Curie  
1867 – 1934



Victor Hess  
1883 – 1964

## Above or Below?

In the late 1800s it was known that gold leaf electroscopes lost their charge due to background radiation ...

...but was it coming from **above** or from **below**?



In 1910 Theodore Wulf took an electroscope to the top of the Eiffel Tower.

The radiation levels decreased by far less than would be expected if the radiation was coming **only** from the ground.


Is the tower radioactive?

Is the atmosphere radioactive?

## Above or Below?

To resolve the question of '... from above or from below?' Hess figured that the Eiffel Tower was not high enough.

He designed detectors that would work at high altitudes and in 1911 and 1912 he took them on a series of balloon flights.



# Cosmic Rays

## Cosmic Radiation


1911 – Ascent to 1000m – radiation levels essentially unchanged

1912 – Ascent to 5000m – radiation levels **increased** dramatically

Was the increase due to radiation from the Sun, or from beyond?

Measurements during a **solar eclipse** effectively ruled out the Sun as the source of cosmic rays.

In 1936 Hess was awarded the Nobel Prize in Physics for his "discovery of cosmic radiation".



17 Apr 1912

## Air Showers




Cosmic ray particles that slam into our atmosphere produce a cascade of secondary particles called an air shower

## Air Showers

High-energy **primary** cosmic ray particles collide with atoms in the upper atmosphere, starting a cascade of **secondary** particles.

At each level of the cascade, new particles are created, each sharing the energy of the particle that created them.

A cascade from one primary cosmic ray particle might result in millions of secondary cosmic ray particles at sea level.




## Air Showers

Many of these new particles are unstable and do not live more than a tiny fraction of a second.


A **muon** is a particle similar to an electron, except that it is much more massive and has no part to play in making atoms.

A muon will not live for much more than a few microseconds, but that is long enough for it to reach sea level.




# Cosmic Rays

## Muon Track in Cloud Chamber



## Applications of Cosmic Ray Muons



If there is any variation in density (of the rock or stone) inside the structure, that will affect the absorption of muons. The process is analogous to using the absorption of x-rays to see inside matter.

## What Are the Primary Particles?

So far we have been looking at **secondary** cosmic ray particles.

What about the **primary** particles – the ones that *create* the showers?

Decades of research has told us about the primary cosmic rays ...


90% are protons ( nuclei of H atoms )



9% are  $\alpha$  particles ( nuclei of He atoms )



1% are nuclei of heavy atoms ( the same atoms as found here within the solar system )



## Detecting Cosmic Rays



*How can we detect cosmic rays either before or after they hit our atmosphere?*



# Cosmic Rays

### Radiation Detectors

What's going on inside the detector?

Radiation

Ionisation

Current → Click


+

-


UNIVERSITY OF LIVERPOOL

33


### Cloud Chambers



Inspired by sightings of the Brocken spectre from the top of Ben Nevis in 1894, Charles Wilson developed chambers for studying cloud formation and optical phenomena in moist air.



In 1911 he perfected the first cloud chamber.




UNIVERSITY OF LIVERPOOL

34



### Cloud Chambers

Wilson realised that when charged particles passed through the cloud chamber water droplets condensed to form visible tracks.

As a result, his cloud chambers had an important role in experimental particle physics for decades.



In 1927 he was awarded the Nobel Prize in Physics for the "most original and wonderful instrument in scientific history".



1911 → 2024

UNIVERSITY OF LIVERPOOL

35

### Cloud Chambers

My cloud chamber is a distant cousin of Wilson's original invention. Details of its design and construction will be left for another day.





UNIVERSITY OF LIVERPOOL

36

# Cosmic Rays

**Demonstration**



Cosmic rays seen in a cloud chamber

UNIVERSITY OF LIVERPOOL


37

**Cosmic Ray Experiments**

Voyager 1 and Voyager 2 carried instruments to measure cosmic rays as they flew through the interplanetary medium.

In 2012 Voyager 1 measured a significant increase in galactic cosmic rays, indicating that it had entered interstellar space.


Travelling about 10% slower, Voyager 2 measured a similar increase in cosmic rays in 2018.



UNIVERSITY OF LIVERPOOL

38


**Cosmic Ray Experiments**



UNIVERSITY OF LIVERPOOL

39

**Cosmic Ray Experiment on the ISS**



UNIVERSITY OF LIVERPOOL

40

# Cosmic Rays

## Cosmic Ray Observatory

There are a number of ground-based cosmic ray observatories. One of the biggest is the Pierre Auger Observatory in Argentina.

To catch a reasonable fraction of the particles created in an air shower the detectors have to be spread over a large area.

For the PAO, this is the area of... Luxembourg.



## Cherenkov Radiation



## Cherenkov Radiation

Development of an air shower




It's blue, and it's spectacular

## Cherenkov Telescope Array





CTA north, La Palma, Canary Islands



CTA south, Atacama Desert, Chile

# Cosmic Rays



## Sources of Cosmic Rays

Where do cosmic rays come from?  
How do they gain so much energy?


UNIVERSITY OF LIVERPOOL

47

### Aside: Spiral Nebulae are Galaxies

**100 years ago** Edwin Hubble reported that ...

*... thirty-six [Cepheid] variable stars were discovered in the two spirals, Andromeda and M33. The study of the periods of these stars at once provided the means of determining the distances of these objects. The results are striking in their confirmation of the view that these spiral nebulae are distant stellar systems. They are found to be at a distance of the order of 1,000,000 light years.*



New York Times, 23 Nov 1924

This was the nail in the coffin of the Great Debate that took place in the early 1920s – "spiral nebulae" are galaxies in their own right.

UNIVERSITY OF LIVERPOOL

48

# Cosmic Rays

## Cosmic Origins

Hess showed that (most) cosmic rays do **not** come from the Sun.

So where **do** they come from?

Sources in the Milky Way? ... or beyond?



Hubble's observations made in the 1920s led to a paradigm shift in our understanding of the Universe ... it is a **lot** bigger than was thought, and is full of billions of galaxies like our own.


So are the sources of cosmic rays galactic, or extra-galactic?

## Supernovae and SMBH




Violent events like supernovae or jets from supermassive black holes produce shock waves in the gas around them.

These shock waves can result in the acceleration of charged particles to **very** high energies.



## Galactic Jets from Hercules A




## Cosmic Sources

Pinning down the sources of cosmic rays?

Shouldn't that be easy?

Measure the direction of incoming cosmic rays and trace their direction back to the source.


Simples!



# Cosmic Rays

## Magnetic Fields


No, not so simple.  
The paths of charged particles are **deflected** in a magnetic field...  
... and magnetic fields are everywhere.



Charged particles spiralling along Earth's magnetic field lines


## Magnetic Fields

Iron filings can reveal the magnet field around a simple bar magnet



## Galactic Magnetic Fields


Dust particles floating around the Milky Way behave like iron filings



Planck


## Galactic Magnetic Fields

Magnetic fields embedded in the Milky Way have a complex structure.  
The fields deflect charged particles and so scramble the direction from which cosmic rays appear to originate.



# Cosmic Rays

### Magnetic Fields in M51




HST + SOFIA

UNIVERSITY OF LIVERPOOL

57

### Inter-Galactic Magnetic Fields







Galactic magnetic fields make it difficult to trace cosmic rays to their source

UNIVERSITY OF LIVERPOOL

58

### Energy Distribution




	x-rays	$\alpha$	$\beta$	$\gamma$				
								
eV	keV	MeV	GeV	TeV	PeV	EeV	ZeV	
$10^0$	$10^3$	$10^6$	$10^9$	$10^{12}$	$10^{15}$	$10^{18}$	$10^{21}$	
Particle Energy			↑ ↑ LHC FCC particle accelerators					

UNIVERSITY OF LIVERPOOL

59

### Energy Distribution



Flux of particles  $m^{-2}sr^{-s}GeV$

←  $1/m^2/s$

$1/m^2/yr$  →

$1/km^2/yr$  →

ultra-high energy CR

Energy / eV

UNIVERSITY OF LIVERPOOL

60

# Cosmic Rays

### Amaterasu Event

Telescope Array  
Utah USA

27 May 2021 APOD 5 Dec 2023

Osaka Metropolitan U / Kyoto U / Ryukyusaku Takehigie

UNIVERSITY OF LIVERPOOL

61

### Amaterasu Event

Telescope Array  
Utah USA

Tiramisu Event

Osaka Metropolitan University  
Kyoto University/Ryukyu

UNIVERSITY OF LIVERPOOL

62

### Ultra-High Energies

Ultra-high-energy cosmic rays should be deflected very little by the magnetic fields that permeate the Milky Way and inter-galactic space.

Hence, tracing back the direction of the Amaterasu event should reveal the source in the sky.

The direction points to ... nothing. No supernova, no SMBH could be identified as the source.

More observations are needed to resolve the mystery of cosmic ray origins.

UNIVERSITY OF LIVERPOOL

63

### Close To the Speed of Light

Just how close to the speed of light are these ultra-high-energy cosmic ray particles travelling?

Expressed as a percentage of the speed of light, the most energetic particle yet observed (the "Oh-My-God particle") was travelling at

**99.99999999999999999999999995 %  $c$**

We can attempt to visualise this speed by imagining a race ...

UNIVERSITY OF LIVERPOOL

64



# Cosmic Rays


### Race With Light

Imagine a 100,000ly race between the most energetic cosmic ray particle and a photon of light.

When the light crosses the finish line, how far behind is the cosmic ray?

After 100,000ly the cosmic ray would be about **5mm** behind.

The time difference would be less than a nanosecond.



The diagram shows a perspective view of a race track on a dark background. The track starts at a 'START' line and ends at a 'FINISH' line. Two parallel paths are shown: a straight path for 'light' and a slightly curved path for 'cosmic ray'. Arrows indicate the direction of travel. The track is set against a background of a galaxy.

UNIVERSITY OF LIVERPOOL

65


### Summary

- Cosmic ray particles are accelerated ... somewhere, somehow
- **Supernovae** and **SuperMassive Black Holes** are probably sources
- Cosmic rays are **deflected** by magnetic fields
  - Lower energy particles get their paths scrambled
  - Higher energy particles travel in straighter lines
- Some particles make it past Earth's protective **magnetic field**
- Hitting the atmosphere creates an **air shower** of secondary particles
- Some of the **secondary** cosmic ray particles make it to sea level
  - These include muons, electrons and protons
  - They make clicks in Geiger counters and tracks in cloud chambers

UNIVERSITY OF LIVERPOOL

66

### Cosmic Rays



The photograph shows a dark, rectangular object, possibly a cloud chamber or bubble chamber, with several bright, white, linear tracks visible. The tracks are of varying lengths and orientations, some appearing as straight lines and others as more complex, branching patterns. The object is set against a dark background.

UNIVERSITY OF LIVERPOOL

67

[www.liverpool.ac.uk/~sdb/Talks](http://www.liverpool.ac.uk/~sdb/Talks)

Dr Steve Barrett

SU3A 22 May 2024



The graphic features a blue and white background with a grid of letters. The letters are arranged in a 5x4 grid, with the words 'STAR WARS HAIL RAIN' and 'COSMIC RAYS' spelled out. The letters are in various colors (yellow, green, grey) and are set against a background of a blue and white grid. The word 'COSMIC' is in white, and 'RAYS' is in green. The background shows a blue and white grid of letters.

COSMIC RAYS